

$\mathbb{Z}$ -actions on AH algebras and $\mathbb{Z}^2$ -actions on AF algebras

Hiroki Matui
 Graduate School of Science
 Chiba University
 Inage-ku, Chiba 263-8522, Japan

Abstract

We consider $\mathbb{Z}$ -actions (single automorphisms) on a unital simple AH algebra with real rank zero and slow dimension growth and show that the uniform outerness implies the Rohlin property under some technical assumptions. Moreover, two $\mathbb{Z}$ -actions with the Rohlin property on such a C^* -algebra are shown to be cocycle conjugate if they are asymptotically unitarily equivalent. We also prove that locally approximately inner and uniformly outer $\mathbb{Z}^2$ -actions on a unital simple AF algebra with a unique trace have the Rohlin property and classify them up to cocycle conjugacy employing the OrderExt group as classification invariants.

1 Introduction

Classification of group actions is one of the most fundamental subjects in the theory of operator algebras. For AFD factors, a complete classification is known for actions of countable amenable groups. However, classification of automorphisms or group actions on C^* -algebras is still a far less developed subject, partly because of K -theoretical difficulties. For AF and AT algebras, A. Kishimoto [9, 10, 11, 12] showed the Rohlin property for a certain class of automorphisms and obtained a cocycle conjugacy result. Following the strategy developed by Kishimoto, H. Nakamura [24] showed that aperiodic automorphisms on unital Kirchberg algebras are completely classified by their KK -classes up to KK -trivial cocycle conjugacy. As for $\mathbb{Z}^N$ -actions, Nakamura [23] introduced the notion of the Rohlin property and classified product type actions of $\mathbb{Z}^2$ on UHF algebras. T. Katsura and the author [8] gave a complete classification of uniformly outer $\mathbb{Z}^2$ -actions on UHF algebras by using the Rohlin property. For Kirchberg algebras, M. Izumi and the author [7] classified a large class of $\mathbb{Z}^2$ -actions and also showed the uniqueness of $\mathbb{Z}^N$ -actions on the Cuntz algebras $\mathcal{O}_2$ and $\mathcal{O}_\infty$. The present article is a continuation of these works.

In the first half of this paper, we study single automorphisms (i.e. $\mathbb{Z}$ -actions) on a unital simple classifiable AH algebra. More precisely, we prove the Rohlin type theorem (Theorem 4.8) for an automorphism α of a unital simple AH algebra A with real rank zero and slow dimension growth under the assumption that α^r is approximately inner for some $r \in \mathbb{N}$ and A has finitely many extremal tracial states. Furthermore, we also prove that if two automorphisms α and β of a unital simple AH algebra with real rank zero and slow dimension growth have the Rohlin property and $\alpha \circ \beta^{-1}$ is asymptotically inner, then the two $\mathbb{Z}$ -actions generated by α and β are cocycle conjugate (Theorem 4.9). These results are generalizations of Kishimoto's work for AF and AT algebras ([9, 10, 11, 12]). For

the proofs, we need to improve some of the arguments in [11, 12] concerning projections and unitaries in central sequence algebras. As a byproduct, it will be also shown that for A as above, the central sequence algebra A_ω satisfies Blackadar's second fundamental comparability question (Proposition 3.8).

In the latter half of the paper, we study $\mathbb{Z}^2$ -actions on a unital simple AF algebra A with a unique trace. We first show a $\mathbb{Z}$ -equivariant version of the Rohlin type theorem for single automorphisms (Theorem 5.5), and as its corollary we obtain the Rohlin type theorem for a $\mathbb{Z}^2$ -action $\varphi : \mathbb{Z}^2 \curvearrowright A$ such that $\varphi_{(r,0)}$ and $\varphi_{(0,s)}$ are approximately inner for some $r, s \in \mathbb{N}$ (Corollary 5.6). This is a generalization of [23, Theorem 3]. Next, by using a $\mathbb{Z}$ -equivariant version of the Evans-Kishimoto intertwining argument [4], we classify uniformly outer locally KK -trivial $\mathbb{Z}^2$ -actions on A up to KK -trivial cocycle conjugacy (Theorem 6.6). This is a generalization of [8, Theorem 6.5]. We remark that KK -triviality of $\alpha \in \text{Aut}(A)$ is equivalent to α being approximately inner, because A is assumed to be AF. For classification invariants, we employ the OrderExt group introduced in [13]. The crossed product of A by the first generator $\varphi_{(1,0)}$ is known to be a unital simple AT algebra with real rank zero. The second generator $\varphi_{(0,1)}$ naturally extends to an automorphism of the crossed product and its OrderExt class gives the invariant of the $\mathbb{Z}^2$ -action φ . However we do not know the precise range of the invariant in general.

2 Preliminaries

We collect notations and terminologies relevant to this paper. For a Lipschitz continuous map f between metric spaces, $\text{Lip}(f)$ denotes the Lipschitz constant of f . Let A be a unital C^* -algebra. For $a, b \in A$, we mean by $[a, b]$ the commutator $ab - ba$. For a unitary $u \in A$, the inner automorphism induced by u is written by $\text{Ad } u$. An automorphism $\alpha \in \text{Aut}(A)$ is called outer, when it is not inner. A single automorphism α is often identified with the $\mathbb{Z}$ -action induced by α . An automorphism α of A is said to be asymptotically inner, if there exists a continuous family of unitaries $\{u_t\}_{t \in [0, \infty)}$ in A such that

$$\alpha(a) = \lim_{t \rightarrow \infty} \text{Ad } u_t(a)$$

for all $a \in A$. When there exists a sequence of unitaries $\{u_n\}_{n \in \mathbb{N}}$ in A such that

$$\alpha(a) = \lim_{n \rightarrow \infty} \text{Ad } u_n(a)$$

for all $a \in A$, α is said to be approximately inner. The set of approximately inner automorphisms of A is denoted by $\overline{\text{Inn}}(A)$. Two automorphisms α and β are said to be asymptotically (resp. approximately) unitarily equivalent if $\alpha \circ \beta^{-1}$ is asymptotically (resp. approximately) inner. The set of tracial states on A is denoted by $T(A)$. We mean by $\text{Aff}(T(A))$ the space of $\mathbb{R}$ -valued affine continuous functions on $T(A)$. The dimension map $D_A : K_0(A) \rightarrow \text{Aff}(T(A))$ is defined by $D_A([p])(\tau) = \tau(p)$. For a homomorphism ρ between C^* -algebras, $K_0(\rho)$ and $K_1(\rho)$ mean the induced homomorphisms on K -groups.

As for group actions on C^* -algebras, we freely use the terminology and notation described in [7, Definition 2.1].

Let A be a separable C^* -algebra and let $\omega \in \beta\mathbb{N} \setminus \mathbb{N}$ be a free ultrafilter. We set

$$c_0(A) = \{(a_n) \in \ell^\infty(\mathbb{N}, A) \mid \lim_{n \rightarrow \infty} \|a_n\| = 0\}, \quad A^\infty = \ell^\infty(\mathbb{N}, A)/c_0(A),$$

$$c_\omega(A) = \{(a_n) \in \ell^\infty(\mathbb{N}, A) \mid \lim_{n \rightarrow \omega} \|a_n\| = 0\}, \quad A^\omega = \ell^\infty(\mathbb{N}, A)/c_\omega(A).$$

We identify A with the C^* -subalgebra of A^∞ (resp. A^ω) consisting of equivalence classes of constant sequences. We let

$$A_\infty = A^\infty \cap A', \quad A_\omega = A^\omega \cap A'$$

and call them the central sequence algebras of A . A sequence $(x_n)_n \in \ell^\infty(\mathbb{N}, A)$ is called a central sequence if $\|[a, x_n]\| \rightarrow 0$ as $n \rightarrow \infty$ for all $a \in A$. A central sequence is a representative of an element in A_∞ . An ω -central sequence is defined in a similar way. When α is an automorphism on A or an action of a discrete group on A , we can consider its natural extension on A^∞ , A^ω , A_∞ and A_ω . We denote it by the same symbol α .

Next, we would like to recall the definition of uniform outerness introduced by Kishimoto and the definition of the Rohlin property of $\mathbb{Z}^N$ -actions introduced by Nakamura.

Definition 2.1 ([9, Definition 1.2]). An automorphism α of a unital C^* -algebra A is said to be uniformly outer if for any $a \in A$, any non-zero projection $p \in A$ and any $\varepsilon > 0$, there exist projections $p_1, p_2, \dots, p_n$ in A such that $p = \sum p_i$ and $\|p_i a \alpha(p_i)\| < \varepsilon$ for all $i = 1, 2, \dots, n$.

We say that an action α of a discrete group on A is uniformly outer if α_g is uniformly outer for every element g of the group other than the identity element.

Let N be a natural number. Let $\xi_1, \xi_2, \dots, \xi_N$ be the canonical basis of $\mathbb{Z}^N$, that is,

$$\xi_i = (0, 0, \dots, 1, \dots, 0, 0),$$

where 1 is in the i -th component. For $m = (m_1, m_2, \dots, m_N)$ and $n = (n_1, n_2, \dots, n_N)$ in $\mathbb{Z}^N$, $m \leq n$ means $m_i \leq n_i$ for all $i = 1, 2, \dots, N$. For $m = (m_1, m_2, \dots, m_N) \in \mathbb{N}^N$, we let

$$m\mathbb{Z}^N = \{(m_1 n_1, m_2 n_2, \dots, m_N n_N) \in \mathbb{Z}^N \mid (n_1, n_2, \dots, n_N) \in \mathbb{Z}^N\}.$$

For simplicity, we denote $\mathbb{Z}^N/m\mathbb{Z}^N$ by $\mathbb{Z}_m$. Moreover, we may identify $\mathbb{Z}_m = \mathbb{Z}^N/m\mathbb{Z}^N$ with

$$\{(n_1, n_2, \dots, n_N) \in \mathbb{Z}^N \mid 0 \leq n_i \leq m_i - 1 \quad \forall i = 1, 2, \dots, N\}.$$

Definition 2.2 ([23, Definition 1]). Let φ be an action of $\mathbb{Z}^N$ on a unital C^* -algebra A . Then φ is said to have the Rohlin property, if for any $m \in \mathbb{N}$ there exist $R \in \mathbb{N}$ and $m^{(1)}, m^{(2)}, \dots, m^{(R)} \in \mathbb{N}^N$ with $m^{(1)}, \dots, m^{(R)} \geq (m, m, \dots, m)$ satisfying the following: For any finite subset F of A and $\varepsilon > 0$, there exists a family of projections

$$e_g^{(r)} \quad (r = 1, 2, \dots, R, \quad g \in \mathbb{Z}_{m^{(r)}})$$

in A such that

$$\sum_{r=1}^R \sum_{g \in \mathbb{Z}_{m^{(r)}}} e_g^{(r)} = 1, \quad \|[a, e_g^{(r)}]\| < \varepsilon, \quad \|\varphi_{\xi_i}(e_g^{(r)}) - e_{g+\xi_i}^{(r)}\| < \varepsilon$$

for any $a \in F$, $r = 1, 2, \dots, R$, $i = 1, 2, \dots, N$ and $g \in \mathbb{Z}_{m^{(r)}}$, where $g + \xi_i$ is understood modulo $m^{(r)}\mathbb{Z}^N$.

It is clear that if $\varphi : \mathbb{Z}^N \curvearrowright A$ has the Rohlin property, then φ is uniformly outer.

We recall the definition of tracial rank zero introduced by H. Lin.

Definition 2.3 ([14, 15]). A unital simple C^* -algebra A is said to have tracial rank zero if for any finite subset $F \subset A$, any $\varepsilon > 0$ and any non-zero positive element $x \in A$ there exists a finite dimensional subalgebra $B \subset A$ with $p = 1_B$ satisfying the following.

- (1) $\|[a, p]\| < \varepsilon$ for all $a \in F$.
- (2) The distance from pap to B is less than ε for all $a \in F$.
- (3) $1 - p$ is Murray-von Neumann equivalent to a projection in xAx .

In [16], H. Lin gave a classification theorem for unital separable simple nuclear C^* -algebras with tracial rank zero which satisfy the UCT. ([16, Theorem 5.2]). Indeed, the class of such C^* -algebras agrees with the class of all unital simple AH algebras with real rank zero and slow dimension growth.

3 Central sequences

Lemma 3.1. *Let A be a unital separable approximately divisible C^* -algebra. Then for any $m \in \mathbb{N}$, there exists a unital embedding of $M_m \oplus M_{m+1}$ into A_∞ .*

Proof. Let $l = m^2 - 1$. For any finite subset $F \subset A$ and $\varepsilon > 0$, there exists a unital finite dimensional subalgebra $B \subset A$ such that every central summand of B is at least $l \times l$ and $\|[a, b]\| < \varepsilon$ for any $a \in F$ and $b \in B$ with $\|b\| \leq 1$. It is easy to find a unital subalgebra C of B such that $C \cong M_m \oplus M_{m+1}$, which completes the proof. $\square$

The following is Lemma 3.6 of [11].

Lemma 3.2. *Let A be a unital simple AT algebra with real rank zero. For any finite subset $F \subset A$ and $\varepsilon > 0$, there exist a finite subset $G \subset A$, $\delta > 0$ and $k \in \mathbb{N}$ such that the following holds. If $p, q \in A$ are projections satisfying $k[p] \leq [q]$, $\|[a, p]\| < \delta$ and $\|[a, q]\| < \delta$ for all $a \in G$, then there exists a partial isometry $v \in A$ such that $v^*v = p$, $vv^* \leq q$ and $\|[a, v]\| < \varepsilon$ for all $a \in F$.*

We generalize the lemma above to AH algebras.

Lemma 3.3. *The above lemma also holds for any unital simple AH algebra with slow dimension growth and real rank zero.*

Proof. Let A be a unital simple AH algebra with slow dimension growth and real rank zero and let Q be the UHF algebra such that $K_0(Q) \cong \mathbb{Q}$. By the classification theorem ([2, 1, 5]), $A \otimes Q$ is a unital simple AT algebra with real rank zero. Let $F \subset A$ be a finite subset and $\varepsilon > 0$. Applying the lemma above to $\{a \otimes 1 \mid a \in F\} \subset A \otimes Q$ and $\varepsilon/2 > 0$, we obtain a finite subset $G \subset A \otimes Q$, $\delta > 0$ and $k \in \mathbb{N}$. We may assume $G = \{a \otimes 1 \mid a \in G_0\} \cup \{1 \otimes b \mid b \in G_1\}$, where G_0 and G_1 are finite subsets of A and Q , respectively. We may further assume that G_0 contains F and δ is less than $\varepsilon/2$. We will prove that G_0 , δ and $4k$ meet the requirement.

Suppose that $p, q \in A$ are non-zero projections satisfying $4k[p] \leq [q]$, $\|[a, p]\| < \delta$ and $\|[a, q]\| < \delta$ for all $a \in G_0$. By Lemma 3.1, $M_2 \oplus M_3$ embeds into A_∞ . Hence there exist a projection r and a partial isometry s such that

$$r \leq q, \quad s^*s = r, \quad ss^* \leq q - r, \quad 4[r] > [q]$$

and

$$\|[a, s]\| < \delta, \quad \|[a, r]\| < \delta \quad \forall a \in G_0.$$

From $4k[p] \leq [q] < 4[r]$, we get $k[p] < [r]$. It follows that there exists a partial isometry $u \in A \otimes Q$ such that $u^*u = p \otimes 1$, $uu^* \leq r \otimes 1$ and $\|[a \otimes 1, u]\| < \varepsilon/2$ for all $a \in F$. We may assume that u belongs to some $A \otimes M_m \subset A \otimes Q$. Put $u = (u_{i,j})_{1 \leq i,j \leq m}$. Define $w = (w_{i,j})_{1 \leq i,j \leq m+1} \in A \otimes M_{m+1}$ by

$$w_{i,j} = \begin{cases} u_{i,j} & \text{if } 1 \leq i, j \leq m \\ su_{i,1} & \text{if } i \neq m+1 \text{ and } j = m+1 \\ 0 & \text{if } i = m+1. \end{cases}$$

It is not so hard to see that $w^*w = p \otimes 1 \in A \otimes M_{m+1}$ and $ww^* \leq q \otimes 1 \in A \otimes M_{m+1}$. Moreover, one has $\|[a \otimes 1, w]\| < \varepsilon$ for all $a \in F$. Let $v = u \oplus w \in A \otimes (M_m \oplus M_{m+1})$. Then $v^*v = p \otimes 1$, $vv^* \leq q \otimes 1$ and $\|[a \otimes 1, v]\| < \varepsilon$ for all $a \in F$.

By [3] (see also [1, 5]), A is approximately divisible. By Lemma 3.1, there exists a unital homomorphism from $M_m \oplus M_{m+1}$ to A_∞ , and so there exists a unital homomorphism π from $A \otimes (M_m \oplus M_{m+1})$ to A^∞ such that $\pi(a \otimes 1) = a$ for $a \in A$. It follows that $\pi(v)^*\pi(v) = p$, $\pi(v)\pi(v)^* \leq q$ and $\|[a, \pi(v)]\| < \varepsilon$ for all $a \in F$. $\square$

Remark 3.4. By using the lemma above and [27, Theorem 4.5], one can show the following. Let A be a unital simple AH algebra with real rank zero and slow dimension growth. Then for any $\alpha \in \text{Aut}(A)$, there exists $\tilde{\alpha} \in \text{Aut}(A)$ such that $\tilde{\alpha}$ has the Rohlin property in the sense of [12, Definition 4.1] and $\tilde{\alpha}$ is asymptotically unitarily equivalent to α .

The following is a well-known fact. We have been unable to find a suitable reference in the literature, so we include a proof for completeness.

Lemma 3.5. *Let A be a unital separable C^* -algebra and let $(p_n)_n$ be a central sequence of projections. For any extremal trace $\tau \in T(A)$, one has*

$$\lim_{n \rightarrow \infty} |\tau(ap_n) - \tau(a)\tau(p_n)| = 0$$

for all $a \in A$.

Proof. First, we deal with the case that there exists $\varepsilon > 0$ such that $\tau(p_n) \geq \varepsilon$ for all $n \in \mathbb{N}$. Consider a sequence of states

$$\varphi_n(a) = \frac{\tau(ap_n)}{\tau(p_n)}$$

on A . Let ψ be an accumulation point of $\{\varphi_n\}_n$. Since $(p_n)_n$ is a central sequence and $\tau(p_n)$ is bounded from below, one can see that ψ is a trace. For any positive element

$a \in A$, it is easy to see $\varphi_n(a) \leq \varepsilon^{-1}\tau(a)$, and so $\psi(a) \leq \varepsilon^{-1}\tau(a)$. Hence, ψ is equal to τ , because τ is extremal. We have shown that any accumulation point of $\{\varphi_n\}$ is τ , which implies φ_n converges to τ . Therefore, $|\tau(ap_n) - \tau(a)\tau(p_n)|$ goes to zero.

Next, we consider the general case. Fix $a \in A$. Take $\varepsilon > 0$ arbitrarily. We would like to show that $|\tau(ap_n) - \tau(a)\tau(p_n)|$ is less than ε for sufficiently large n . We may assume $\|a\| \leq 1$. Put

$$C = \{n \in \mathbb{N} \mid \tau(p_n) \geq \varepsilon/2\}.$$

If $n \notin C$, then evidently $|\tau(ap_n) - \tau(a)\tau(p_n)|$ is less than ε . By applying the first part of the proof to $(p_n)_{n \in C}$, we have $|\tau(ap_n) - \tau(a)\tau(p_n)| < \varepsilon$ for sufficiently large $n \in C$, thereby completing the proof. $\square$

Lemma 3.6. *Let A be a unital simple separable C^* -algebra with tracial rank zero. Suppose that A has finitely many extremal traces. For any finite subset $F \subset A$ and $\varepsilon > 0$, there exist a finite subset $G \subset A$ and $\delta > 0$ such that the following hold. If $p, q \in A$ are projections satisfying*

$$\|[a, p]\| < \delta, \quad \|[a, q]\| < \delta \quad \forall a \in G$$

*and $\tau(p) + \varepsilon < \tau(q)$ for all $\tau \in T(A)$, then there exists a partial isometry $v \in A$ such that $v^*v \leq p$, $vv^* \leq q$,*

$$\|[a, v]\| < \varepsilon \quad \forall a \in F$$

*and $\tau(p - v^*v) < \varepsilon$ for all $\tau \in T(A)$.*

Proof. The proof is by contradiction. Suppose that the assertion does not hold for a finite subset $F \subset A$ and $\varepsilon > 0$. We would have central sequences of projections $(p_n)_n$ and $(q_n)_n$ such that

$$\tau(p_n) + \varepsilon < \tau(q_n) \quad \forall \tau \in T(A), \quad n \in \mathbb{N}$$

and any partial isometry does not meet the requirement.

Use tracial rank zero to find a projection $e \in A$ and a finite dimensional unital subalgebra $E \subset eAe$ such that the following are satisfied.

- For any $a \in F$, $\|[a, e]\| < \varepsilon/4$.
- For any $a \in F$ there exists $b \in E$ such that $\|eae - b\| < \varepsilon/4$.
- $\tau(1 - e) < \varepsilon$ for all $\tau \in T(A)$.

Since $(p_n)_n$ and $(q_n)_n$ are central sequences of projections, we can find projections p'_n and q'_n in $A \cap E'$ such that $\|p_n - p'_n\| \rightarrow 0$ and $\|q_n - q'_n\| \rightarrow 0$ as $n \rightarrow \infty$. We would like to show that, for sufficiently large n , there exists a partial isometry $v_n \in eAe \cap E'$ such that $v_n^*v_n = ep'_n$ and $v_nv_n^* \leq eq'_n$. Let $\{e_1, e_2, \dots, e_m\}$ be a family of minimal central projections in $eAe \cap E'$ such that $e_1 + e_2 + \dots + e_m = e$. Clearly $e_i(eAe \cap E')$ is a unital simple C^* -algebra with tracial rank zero and the space of tracial states on $e_i(eAe \cap E')$ is naturally identified with $T(A)$. By lemma 3.5, for sufficiently large n , one has $\tau(e_ip'_n) < \tau(e_iq'_n)$ for all $\tau \in T(A)$ and $i = 1, 2, \dots, m$, because A has finitely many extremal traces. Hence $[e_ip'_n] \leq [e_iq'_n]$ in $K_0(e_i(eAe \cap E'))$. It follows that there exists a partial isometry $v_n \in eAe \cap E'$ such that $v_n^*v_n = ep'_n$ and $v_nv_n^* \leq eq'_n$. Besides, $\tau(p'_n - v_n^*v_n) = \tau(p'_n(1 - e)) < \varepsilon$ and $\|[a, v_n]\| < \varepsilon$ for all $a \in F$. This is a contradiction. $\square$

By using Lemma 3.6 and 3.3, we can show the following.

Lemma 3.7. *Let A be a unital simple AH algebra with slow dimension growth and real rank zero. Suppose that A has finitely many extremal traces. For any finite subset $F \subset A$ and $\varepsilon > 0$, there exist a finite subset $G \subset A$ and $\delta > 0$ such that the following hold. If $p, q \in A$ are projections satisfying*

$$\|[a, p]\| < \delta, \quad \|[a, q]\| < \delta \quad \forall a \in G$$

*and $\tau(p) + \varepsilon < \tau(q)$ for all $\tau \in T(A)$, then there exists a partial isometry $u \in A$ such that $u^*u = p$, $uu^* \leq q$ and $\|[a, u]\| < \varepsilon$ for all $a \in F$.*

Proof. Notice that A has tracial rank zero by [15, Proposition 2.6]. Suppose that a finite subset $F \subset A$ and $\varepsilon > 0$ are given. By applying Lemma 3.3 to F and $\varepsilon/2$, we obtain a finite subset $F_1 \subset A$, $\varepsilon_1 > 0$ and $k \in \mathbb{N}$. By applying Lemma 3.6 to $F \cup F_1 \cup F_1^*$ and $\min\{\varepsilon_1/4, \varepsilon/k, \varepsilon/2\}$, we obtain a finite subset $G \subset A$, $\delta > 0$. We would like to show that $G \cup F_1$ and $\min\{\delta, \varepsilon_1/2\}$ meet the requirement. Suppose that $p, q \in A$ are projections satisfying

$$\|[a, p]\| < \min\{\delta, \varepsilon_1/2\}, \quad \|[a, q]\| < \min\{\delta, \varepsilon_1/2\} \quad \forall a \in G \cup F_1$$

and $\tau(p) + \varepsilon < \tau(q)$ for all $\tau \in T(A)$. By Lemma 3.6, there exists a partial isometry $v \in A$ such that $v^*v \leq p$, $vv^* \leq q$,

$$\|[a, v]\| < \min\{\varepsilon_1/4, \varepsilon/2\} \quad \forall a \in F \cup F_1 \cup F_1^*$$

and $\tau(p - v^*v) < \varepsilon/k$ for all $\tau \in T(A)$. Let $p' = p - v^*v$ and $q' = q - vv^*$. One has

$$\tau(q') = \tau(q - vv^*) = \tau(q) - \tau(p) + \tau(p - v^*v) > \varepsilon,$$

and so $k[p'] \leq [q']$. One also has $\|[a, p']\| < \varepsilon_1$ and $\|[a, q']\| < \varepsilon_1$ for all $a \in F_1$. By Lemma 3.3, we obtain a partial isometry $w \in A$ such that $w^*w = p'$, $ww^* \leq q'$ and $\|[a, w]\| < \varepsilon/2$ for all $a \in F$. Put $u = v + w$. It is easy to see $u^*u = p$, $uu^* \leq q$ and $\|[a, u]\| < \varepsilon$ for all $a \in F$. $\square$

Any tracial state $\tau \in T(A)$ naturally extends to a tracial state on A^ω and we write it by τ_ω .

Proposition 3.8. *Let A be a unital simple AH algebra with slow dimension growth and real rank zero. Suppose that A has finitely many extremal traces. If $p, q \in A_\omega$ are projections satisfying $\tau_\omega(p) < \tau_\omega(q)$ for all $\tau \in T(A)$, then there exists $v \in A_\omega$ such that $v^*v = p$ and $vv^* \leq q$. In particular, A_ω satisfies Blackadar's second fundamental comparability question.*

Proof. Let $(p_n)_n$ and $(q_n)_n$ be ω -central sequences of projections such that

$$\lim_{n \rightarrow \omega} \tau(p_n) < \lim_{n \rightarrow \omega} \tau(q_n)$$

for all $\tau \in T(A)$. Since A has finitely many extremal traces, there exists $\varepsilon > 0$ such that

$$C = \{n \in \mathbb{N} \mid \tau(p_n) + \varepsilon < \tau(q_n) \text{ for all } \tau \in T(A)\} \in \omega.$$

We choose an increasing sequence $\{F_m\}_{m=1}^\infty$ of finite subsets of A whose union is dense in A . By applying Lemma 3.7 to F_m and ε/m , we obtain a finite subset $G_m \subset A$ and $\delta_m > 0$. We may assume that $\{G_m\}_m$ is increasing and $\{\delta_m\}_m$ is decreasing. Put

$$C_m = \{n \in C \mid \|[a, p_n]\| < \delta_m \text{ and } \|[a, q_n]\| < \delta_m \text{ for all } a \in G_m\} \in \omega.$$

For $n \in C_m \setminus C_{m+1}$, by Lemma 3.7, there exists a partial isometry u_n such that $u_n^* u_n = p_n$, $u_n u_n^* \leq q_n$ and $\|[a, u_n]\| < \varepsilon/m$ for all $a \in F_m$. For $n \in \mathbb{N} \setminus C_1$, we let $u_n = 0$. Then $(u_n)_n$ is a desired ω -central sequence of partial isometries. $\square$

The following is Lemma 4.4 of [12].

Lemma 3.9. *Let A be a unital simple AT algebra with real rank zero. For any finite subset $F \subset A$ and $\varepsilon > 0$, there exist a finite subset $G \subset A$ and $\delta > 0$ such that the following holds. If $u : [0, 1] \rightarrow A$ is a path of unitaries satisfying $\|[a, u(t)]\| < \delta$ for all $a \in G$ and $t \in [0, 1]$, then there exists a path of unitaries $v : [0, 1] \rightarrow A$ satisfying*

$$v(0) = u(0), \quad v(1) = u(1), \quad \|[a, v(t)]\| < \varepsilon \quad \forall a \in F, \quad t \in [0, 1]$$

and $\text{Lip}(v) < 5\pi + 1$.

We generalize the lemma above to AH algebras.

Lemma 3.10. *The above lemma also holds for any unital simple AH algebra with slow dimension growth and real rank zero, the Lipschitz constant being bounded by 11π .*

Proof. Let A be a unital simple AH algebra with slow dimension growth and real rank zero and let Q be the UHF algebra such that $K_0(Q) \cong \mathbb{Q}$. By the classification theorem ([2, 1, 5]), $A \otimes Q$ is a unital simple AT algebra with real rank zero. Let $F \subset A$ be a finite subset and $\varepsilon > 0$. We may assume that F is contained in the unit ball of A . Applying the lemma above to $\{a \otimes 1 \mid a \in F\} \subset A \otimes Q$ and $\varepsilon/2 > 0$, we obtain a finite subset $G \subset A \otimes Q$ and $\delta > 0$. We may assume $G = \{a \otimes 1 \mid a \in G_0\} \cup \{1 \otimes b \mid b \in G_1\}$, where G_0 and G_1 are finite subsets of A and Q , respectively. We will prove that G_0 and δ meet the requirement.

Suppose that $u : [0, 1] \rightarrow A$ is a path of unitaries satisfying $\|[a, u(t)]\| < \delta$ for all $a \in G_0$ and $t \in [0, 1]$. Choose $N \in \mathbb{N}$ so that $\|u(\frac{k}{N}) - u(\frac{k+1}{N})\| < \varepsilon/2$ for every $k = 0, 1, \dots, N-1$. Put $u_k = u(k/N)$. By the lemma above, we can find continuous paths $x : [0, 1] \rightarrow A \otimes Q$, $y_k : [0, 1] \rightarrow A \otimes Q$ and $z_k : [0, 1] \rightarrow A \otimes Q$ for $k = 1, 2, \dots, N-1$ such that

$$x(0) = u_0 \otimes 1, \quad x(1) = u_N \otimes 1, \quad \|[a \otimes 1, x(t)]\| < \varepsilon/2 \quad \forall a \in F, \quad t \in [0, 1],$$

$$y_k(0) = u_0 \otimes 1, \quad y_k(1) = u_k \otimes 1, \quad \|[a \otimes 1, y_k(t)]\| < \varepsilon/2 \quad \forall a \in F, \quad t \in [0, 1],$$

$$z_k(0) = u_k \otimes 1, \quad z_k(1) = u_N \otimes 1, \quad \|[a \otimes 1, z_k(t)]\| < \varepsilon/2 \quad \forall a \in F, \quad t \in [0, 1],$$

and $\text{Lip}(x), \text{Lip}(y_k), \text{Lip}(z_k)$ are less than $5\pi + 1$. We may assume that the ranges of x, y_k, z_k are contained in $A \otimes M_n$ for some $M_n \subset Q$.

Put $m = n(N-1)$. We would like to construct a path of unitaries $v : [0, 1] \rightarrow A \otimes (M_m \oplus M_{m+1})$ such that $\text{Lip}(v) < 11\pi$, $v(0) = u_0 \otimes 1$, $v(1) = u_N \otimes 1$ and $\|[a \otimes 1, v(t)]\| < \varepsilon$ for all $a \in F$ and $t \in [0, 1]$. First, let $\tilde{x} : [0, 1] \rightarrow A \otimes M_m$ be the direct sum of $N-1$

copies of $x : [0, 1] \rightarrow A \otimes M_n$. Next, by using $y_1, y_2, \dots, y_{N-1}$, we can find a path $\tilde{y} : [0, 1] \rightarrow A \otimes M_{m+1}$ such that

$$\begin{aligned}\tilde{y}(0) &= u_0 \otimes 1, \\ \tilde{y}(1) &= \text{diag}(u_0, \underbrace{u_1, \dots, u_1}_n, \underbrace{u_2, \dots, u_2}_n, \dots, \underbrace{u_{N-1}, \dots, u_{N-1}}_n) \\ \|[a \otimes 1, \tilde{y}(t)]\| &< \varepsilon/2 \quad \forall a \in F, t \in [0, 1]\end{aligned}$$

and $\text{Lip}(\tilde{y}) < 5\pi + 1$. Likewise, by using $z_1, z_2, \dots, z_{N-1}$, we can find a path $\tilde{z} : [0, 1] \rightarrow A \otimes M_{m+1}$ such that

$$\begin{aligned}\tilde{z}(0) &= \text{diag}(\underbrace{u_1, \dots, u_1}_n, \underbrace{u_2, \dots, u_2}_n, \dots, \underbrace{u_{N-1}, \dots, u_{N-1}}_n, u_N) \\ \tilde{z}(1) &= u_N \otimes 1, \\ \|[a \otimes 1, \tilde{z}(t)]\| &< \varepsilon/2 \quad \forall a \in F, t \in [0, 1]\end{aligned}$$

and $\text{Lip}(\tilde{z}) < 5\pi + 1$. Since $\|\tilde{y}(1) - \tilde{z}(0)\| < \varepsilon/2$, if ε is sufficiently small, there exists a path $w : [0, 1] \rightarrow M_{m+1}$ such that

$$w(0) = u_0 \otimes 1, \quad w(1) = u_N \otimes 1, \quad \|[a \otimes 1, w(t)]\| < \varepsilon \quad \forall a \in F, t \in [0, 1],$$

and $\text{Lip}(w) < 11\pi$. Then $v = \tilde{x} \oplus w$ is the desired path.

By [3] (see also [1, 5]), A is approximately divisible. By Lemma 3.1, there exists a unital homomorphism from $M_m \oplus M_{m+1}$ to A_∞ , and so there exists a unital homomorphism π from $A \otimes (M_m \oplus M_{m+1})$ to A^∞ such that $\pi(a \otimes 1) = a$ for $a \in A$. It follows that the path $\tilde{v} : [0, 1] \ni t \mapsto \pi(v(t)) \in A^\infty$ satisfies

$$\tilde{v}(0) = u_0, \quad \tilde{v}(1) = u_N, \quad \|[a, \tilde{v}(t)]\| < \varepsilon \quad \forall a \in F, t \in [0, 1]$$

and $\text{Lip}(\tilde{v}) < 11\pi$, which completes the proof. $\square$

4 Automorphisms of AH algebras

In this section, we discuss the Rohlin property of automorphisms of AH algebras. For $a \in A$, we define

$$\|a\|_2 = \sup_{\tau \in T(A)} \tau(a^*a)^{1/2}.$$

If A is simple and $T(A)$ is non-empty, then $\|\cdot\|_2$ is a norm.

Proposition 4.1. *Let A be a unital simple separable C^* -algebra with tracial rank zero and let $\Gamma \subset \text{Aut}(A)$ be a finite subset containing the identity. Suppose that there exists a sequence of projections $(e_n)_n$ in A satisfying the following property.*

- (1) $\|\gamma(e_n)\gamma'(e_n)\|_2 \rightarrow 0$ for any $\gamma, \gamma' \in \Gamma$ such that $\gamma \neq \gamma'$.
- (2) $\|1 - \sum_{\gamma \in \Gamma} \gamma(e_n)\|_2 \rightarrow 0$.
- (3) For every $a \in A$, we have $\|[a, e_n]\|_2 \rightarrow 0$.

Then there exists a sequence of projections $(f_n)_n$ in A satisfying the following.

- (1) $\|\gamma(f_n)\gamma'(f_n)\| \rightarrow 0$ for any $\gamma, \gamma' \in \Gamma$ such that $\gamma \neq \gamma'$.
- (2) $\|e_n - f_n\|_2 \rightarrow 0$.
- (3) For every $a \in A$, we have $\|[a, f_n]\| \rightarrow 0$.

Proof. This is almost the same as [19, Proposition 5.4]. In [19, Proposition 5.4], the finite set Γ is assumed to be an orbit of a single automorphism γ of finite order. The proof, however, does not need this. $\square$

The following is a variant of [9, Lemma 3.1] and [26, Theorem 2.17]. See [26, Definition 2.1] for the definition of the tracial Rohlin property.

Theorem 4.2. *Let A be a unital simple separable C^* -algebra with tracial rank zero. Suppose that A has finitely many extremal traces. Let α be an automorphism of A such that α^m is uniformly outer for any $m \in \mathbb{N}$. Then α has the tracial Rohlin property.*

Proof. Let $\{\tau_1, \dots, \tau_d\}$ be the set of extremal tracial states of A and let (π_i, H_i) be the GNS representation associated with τ_i . It is well-known that $\pi_i(A)''$ is a hyperfinite II_1 -factor (see [26, Lemma 2.16]). Let $\rho = \bigoplus_{i=1}^d \pi_i$. Note that, for a bounded sequence $(a_n)_n$ in A , $\rho(a_n)$ converges to zero in the strong operator topology if and only if $\|a_n\|_2$ converges to zero. We regard A as a subalgebra of $N = \rho(A)'' \cong \bigoplus_{i=1}^d \pi_i(A)''$ and denote the extension of the automorphism α to N by $\bar{\alpha}$. Let k be the minimum positive integer such that $\tau_i \circ \alpha^k = \tau_i$ for all $i = 1, 2, \dots, d$. In the same way as [9, Lemma 3.1], for any $l \in \mathbb{N}$, one can find a sequence $\{f_0^{(j)}, \dots, f_{kl-1}^{(j)}\}$ of orthogonal families of projections in N such that $\sum_{i=0}^{kl-1} f_i^{(j)} = 1$,

$$\begin{aligned} [a, f_i^{(j)}] &\rightarrow 0 \quad \forall a \in A, \\ \bar{\alpha}(f_i^{(j)}) - f_{i+1}^{(j)} &\rightarrow 0 \quad \forall i = 0, 1, \dots, kl-1 \end{aligned}$$

in the strong operator topology as $j \rightarrow \infty$, where $f_{kl}^{(j)} = f_0^{(j)}$. By [26, Lemma 2.15], we may replace the projections $f_i^{(j)}$ with projections of A . From the proposition above, we can conclude that α has the tracial Rohlin property. $\square$

The following is a well-known fact, but we include the proof for the reader's convenience.

Lemma 4.3. *Let A be a unital separable C^* -algebra and let $\alpha \in \overline{\text{Inn}}(A)$. For any separable subset $C \subset A_\infty$, there exists a unitary $u \in A_\infty$ such that $uxu^* = \alpha(x)$ for all $x \in C$.*

Proof. We choose an increasing sequence $\{F_n\}_{n \in \mathbb{N}}$ of finite subsets of A whose union is dense in A . We can find a sequence of unitaries $(v_n)_n$ in A such that

$$\|v_n a v_n^* - \alpha^{-1}(a)\| < n^{-1}$$

for all $a \in F_n$, because α is approximately inner. We may assume that C is countable. Let $C = \{x_1, x_2, \dots\}$ and let $(x_{i,j})_j$ be a representative of x_i . There exists an increasing sequence $(m(n))_n$ of natural numbers such that

$$\|[v_n, x_{i,j}]\| < n^{-1} \quad \forall j \geq m(n)$$

for any $i = 1, 2, \dots, n$, because $(x_{i,j})_j$ is a central sequence. Since α is in $\overline{\text{Inn}}(A)$, one can find a sequence of unitaries $(w_n)_n$ in A such that

$$\|w_n a w_n^* - \alpha(a)\| < n^{-1}$$

for all a in

$$\alpha^{-1}(F_n) \cup \{x_{i,j} \mid i = 1, \dots, n, m(n) \leq j < m(n+1)\}.$$

For $j \in \mathbb{N}$, find $n \in \mathbb{N}$ so that $m(n) \leq j < m(n+1)$ and define a unitary u_j by $u_j = w_n v_n$. It is easy to see

$$\|[u_j, a]\| < 2/n \quad \forall a \in F_n$$

and

$$\|u_j x_{i,j} u_j^* - \alpha(x_{i,j})\| < 2/n \quad \forall i = 1, \dots, n,$$

and so the proof is completed. $\square$

We quote the following theorem by Lin and Osaka from [20]. See [20, Definition 2.4] for the definition of the tracial cyclic Rohlin property. Since we need to discuss an ‘equivariant version’ of this theorem later, we would like to include the proof briefly.

Theorem 4.4 ([20, Theorem 3.4]). *Let A be a unital simple separable C^* -algebra and suppose that the order on projections in A is determined by traces. Suppose that $\alpha \in \text{Aut}(A)$ has the tracial Rohlin property. If α^r is in $\overline{\text{Inn}}(A)$ for some $r \in \mathbb{N}$, then α has the tracial cyclic Rohlin property.*

Proof. Take $m \in \mathbb{N}$ and $\varepsilon > 0$ arbitrarily. Let $l = r(mr + 1)$. Since α has the tracial Rohlin property and the order on projections is determined by traces, there exists a central sequence of projections $(e_n)_n$ such that

$$\lim_{n \rightarrow \infty} \|e_n \alpha^i(e_n)\| = 0 \quad \forall i = 1, 2, \dots, l-1$$

and

$$\lim_{n \rightarrow \infty} \tau(1 - (e_n + \alpha(e_n) + \dots + \alpha^{l-1}(e_n))) = 0 \quad \forall \tau \in T(A).$$

Let $e \in A_\infty$ be the image of $(e_n)_n$ and define $\tilde{e}$ by

$$\tilde{e} = \sum_{i=0}^{r-1} \alpha^{i(mr+1)}(e).$$

It follows from the lemma above that there exists a partial isometry $v \in A_\infty$ such that $v^*v = \tilde{e}$ and $vv^* = \alpha(\tilde{e})$. The C^* -algebra C generated by $v, \alpha(v), \dots, \alpha^{mr-1}(v)$ is isomorphic to M_{mr+1} and its unit is $\tilde{e} + \alpha(\tilde{e}) + \dots + \alpha^{mr}(\tilde{e})$. The rest of the proof is exactly the same as that of [9, Lemma 4.3] and we omit it. $\square$

Remark 4.5. The following was shown by Lin in [17, Theorem 3.4]. Let A be a unital simple separable C^* -algebra with tracial rank zero. Suppose that $\alpha \in \text{Aut}(A)$ has the tracial cyclic Rohlin property and that there exists $r \in \mathbb{N}$ such that $K_0(\alpha^r)|_G = \text{id}_G$ for some subgroup $G \subset K_0(A)$ for which $D_A(G)$ is dense in $D_A(K_0(A))$. Then $A \rtimes_\alpha \mathbb{Z}$ has tracial rank zero.

By using Lemma 3.3, we can show the following.

Lemma 4.6. *Let A be a unital simple AH algebra with slow dimension growth and real rank zero. Suppose that $\alpha \in \text{Aut}(A)$ has the tracial cyclic Rohlin property and that there exists $r \in \mathbb{N}$ such that $\tau \circ \alpha^r = \tau$ for any $\tau \in T(A)$. Then, for any $m \in \mathbb{N}$, there exist projections $e, f \in A_\infty$ and a partial isometry $v \in A_\infty$ such that*

$$v^*v = f, \quad vv^* \leq e, \quad f + \sum_{i=0}^{mr} \alpha^i(e) = 1$$

and $\alpha^{mr+1}(e) = e$.

Proof. Suppose that we are given $m \in \mathbb{N}$. Let $l = r(mr+1)$. Since α has the tracial cyclic Rohlin property, we can find central sequences of projections $(e_n)_n$ and $(f_n)_n$ such that

$$f_n + \sum_{i=0}^{l-1} \alpha^i(e_n) \rightarrow 1, \quad e_n - \alpha^l(e_n) \rightarrow 0, \quad \sup_{\tau \in T(A)} \tau(f_n) \rightarrow 0$$

as $n \rightarrow \infty$. There exists a central sequence of projections $(\tilde{e}_n)_n$ such that

$$\lim_{n \rightarrow \infty} \tilde{e}_n - \sum_{i=0}^{r-1} \alpha^{i(mr+1)}(e_n) = 0.$$

Then

$$f_n + \sum_{i=0}^{mr} \alpha^i(\tilde{e}_n) \rightarrow 1, \quad \tilde{e}_n - \alpha^{mr+1}(\tilde{e}_n) \rightarrow 0$$

as $n \rightarrow \infty$. It is also easy to see $\tau(\alpha(\tilde{e}_n)) = \tau(\tilde{e}_n)$ for all $\tau \in T(A)$, and so $\tau(\tilde{e}_n)$ goes to $(mr+1)^{-1}$ for all $\tau \in T(A)$. Therefore, by Lemma 3.3, one can find a central sequence of partial isometries $(v_n)_n$ such that $v_n^*v_n = f_n$ and $v_nv_n^* \leq \tilde{e}_n$ for sufficiently large n , which completes the proof. $\square$

By using the lemma above, we can show the following theorem.

Theorem 4.7. *Let A be a unital simple AH algebra with slow dimension growth and real rank zero. Suppose that $\alpha \in \text{Aut}(A)$ has the tracial cyclic Rohlin property. If there exists $r \in \mathbb{N}$ such that $\tau \circ \alpha^r = \tau$ for any $\tau \in T(A)$, then α has the Rohlin property.*

Proof. Suppose that we are given $M \in \mathbb{N}$. Choose a natural number $m \in \mathbb{N}$ so that $m \geq M$ and $m \equiv 1 \pmod{r}$. Let k, l be sufficiently large natural numbers satisfying $k \equiv l \equiv 1 \pmod{r}$. By the lemma above, we can find projections $e, f \in A_\infty$ and a partial isometry $v \in A_\infty$ such that

$$v^*v = f, \quad vv^* \leq e, \quad f + \sum_{i=0}^{klm-1} \alpha^i(e) = 1$$

and $\alpha^{klm}(e) = e$. Define $\tilde{e}, w \in A_\infty$ by

$$\tilde{e} = \sum_{i=0}^{k-1} \alpha^{ilm}(e) \quad \text{and} \quad w = \frac{1}{\sqrt{k}} \sum_{i=0}^{k-1} \alpha^{ilm}(v).$$

Then $\tilde{e}$ is a projection and w is a partial isometry satisfying

$$f + \sum_{i=0}^{lm-1} \alpha^i(\tilde{e}) = 1, \quad \alpha^{lm}(\tilde{e}) = \tilde{e}$$

and

$$w^*w = f, \quad ww^* \leq \tilde{e}, \quad \|\alpha^{lm}(w) - w\| \leq \frac{2}{\sqrt{k}}.$$

Let D be the C^* -algebra generated by $w, \alpha(w), \dots, \alpha^{lm-1}(w)$. Then D is isomorphic to M_{lm+1} and the unit 1_D of D is equal to $f + ww^* + \dots + \alpha^{lm-1}(ww^*)$. From the spectral property of α restricted to D , if k and l are sufficiently large, we can obtain projections $p_0, \dots, p_{m-1}, q_0, \dots, q_m$ of D such that

$$\sum_{i=1}^{m-1} p_i + \sum_{i=1}^m q_i = 1_D, \quad \alpha(p_i) \approx p_{i+1}, \quad \alpha(q_i) \approx q_{i+1},$$

where $p_m = p_0$ and $q_{m+1} = q_0$. We define projections p'_i in A_∞ by

$$p'_i = p_i + \sum_{j=0}^{l-1} \alpha^{i+jm}(\tilde{e} - ww^*).$$

Then the projections $p'_0, \dots, p'_{m-1}, q_0, \dots, q_m$ meet the requirement. See [9, 10] for details. $\square$

Combining the theorems above, we obtain the following theorem which is a generalization of [11, Theorem 2.1].

Theorem 4.8. *Let A be a unital simple AH algebra with slow dimension growth and real rank zero and let $\alpha \in \text{Aut}(A)$. Suppose that A has finitely many extremal traces and that α^r is approximately inner for some $r \in \mathbb{N}$. Then the following are equivalent.*

- (1) α has the Rohlin property.
- (2) α^m is uniformly outer for any $m \in \mathbb{N}$.

We can also generalize [12, Theorem 5.1] by using Lemma 3.10 instead of [12, Lemma 4.4].

Theorem 4.9. *Let A be a unital simple AH algebra with slow dimension growth and real rank zero. If $\alpha, \beta \in \text{Aut}(A)$ have the Rohlin property and α is asymptotically unitarily equivalent to β , then there exist $\mu \in \overline{\text{Inn}}(A)$ and a unitary $u \in A$ such that*

$$\text{Ad } u \circ \alpha = \mu \circ \beta \circ \mu^{-1}.$$

The proof is similar to that of [12, Theorem 5.1] and we omit it.

As an application of the theorems above, we can show the following, which will be used in Section 6.

Lemma 4.10. *Let A be a unital simple AF algebra with finitely many extremal traces and let α be an approximately inner automorphism of A such that α^m is uniformly outer for all $m \in \mathbb{N}$. For any finite subset $F \subset A$ and $\varepsilon > 0$, there exist a finite subset $G \subset A$ and $\delta > 0$ satisfying the following. If $u : [0, 1] \rightarrow A$ is a path of unitaries such that*

$$\|[a, u(t)]\| < \delta \quad \text{and} \quad \|u(t) - \alpha(u(t))\| < \delta \quad \forall a \in G, \quad t \in [0, 1],$$

then there exists a path of unitaries $v : [0, 1] \rightarrow A$ such that

$$v(0) = u(0), \quad v(1) = u(1), \quad \|[a, v(t)]\| < \varepsilon, \quad \|v(t) - \alpha(v(t))\| < \varepsilon, \quad \forall a \in F, \quad t \in [0, 1]$$

and $\text{Lip}(v) < 2\pi$.

Proof. Let x_n be a unitary of $M_n(\mathbb{C})$ such that $\text{Sp}(x_n) = \{\omega^k \mid k = 0, 1, \dots, n-1\}$, where $\omega = \exp(2\pi\sqrt{-1}/n)$. One can find an increasing sequence $\{A_n\}_{n=1}^\infty$ of unital finite dimensional subalgebras of A such that $\bigcup_n A_n$ is dense in A and there exists a unital embedding $\pi_n : M_n \oplus M_{n+1} \rightarrow A_{n+1} \cap A'_n$. Let $y_n = \pi_n(x_n \oplus x_{n+1})$. Define an automorphism σ of A by $\sigma = \lim_{n \rightarrow \infty} \text{Ad}(y_1 y_2 \dots y_n)$. Then σ is approximately inner and σ^m is uniformly outer for all $m \in \mathbb{N}$.

We would like to show that the assertion holds for σ . Suppose that we are given $F \subset A$ and $\varepsilon > 0$. Without loss of generality, we may assume that there exists $n \in \mathbb{N}$ such that F is contained in the unit ball of A_n . Applying [8, Lemma 4.2] to $\varepsilon/2$, we obtain a positive real number $\delta_1 > 0$. We may assume δ_1 is less than $\min\{2, \varepsilon\}$. Choose a finite subset $G \subset A$ and $\delta_2 > 0$ so that if $z : [0, 1] \rightarrow A$ is a path of unitaries such that $\|[a, z(t)]\| < \delta_2$ for all $a \in G$ and $t \in [0, 1]$, then there exists a path of unitaries $\tilde{z} : [0, 1] \rightarrow A \cap A'_n$ such that $\|z(t) - \tilde{z}(t)\| < \delta_1/6$. Let $\delta = \min\{\delta_1/6, \delta_2\}$. Suppose that $u : [0, 1] \rightarrow A$ is a path of unitaries such that

$$\|[a, u(t)]\| < \delta \quad \text{and} \quad \|u(t) - \sigma(u(t))\| < \delta \quad \forall a \in G, \quad t \in [0, 1].$$

By the choice of δ , we can find $\tilde{u} : [0, 1] \rightarrow A \cap A'_n$ such that $\|u(t) - \tilde{u}(t)\| < \delta_1/6$. We may assume that there exists $m > n$ such that the range of $\tilde{u}$ is contained in A_m . Put $y = y_n y_{n+1} \dots y_{m-1} \in A_m \cap A'_n$. Then

$$\|[y, \tilde{u}(t)]\| = \|\tilde{u}(t) - \sigma(\tilde{u}(t))\| < \|u(t) - \sigma(u(t))\| + \delta_1/3 < \delta + \delta_1/3 \leq \delta_1/2$$

for every $t \in [0, 1]$. Hence $\|[y, \tilde{u}(t)\tilde{u}(0)^*]\|$ is less than δ_1 . It follows from [8, Lemma 4.2] that one can find a path of unitaries $w : [0, 1] \rightarrow A_m \cap A'_n$ such that

$$w(0) = 1, \quad w(1) = \tilde{u}(1)\tilde{u}(0)^*, \quad \text{Lip}(w) \leq \pi + \varepsilon$$

and $\|[y, w(t)]\| < \varepsilon/2$ for every $t \in [0, 1]$. Note that $yw(t)y^*$ is equal to $\sigma(w(t))$. By perturbing $w(t)u(0)$ a little bit, the required $v : [0, 1] \rightarrow A$ is obtained.

Suppose that α is an approximately inner automorphism of A such that α^m is uniformly outer for all $m \in \mathbb{N}$. By Theorem 4.8 and Theorem 4.9, there exist $\mu \in \text{Aut}(A)$ and a unitary $u \in A$ such that $\text{Ad } u \circ \alpha = \mu \circ \sigma \circ \mu^{-1}$. Moreover, one can choose u arbitrarily close to 1, because A is AF (see [6, 12]). Therefore the assertion also holds for α . $\square$

Remark 4.11. In the proof of the lemma above, it is easily seen that (the $\mathbb{Z}$ -action generated by) σ is asymptotically representable ([7, Definition 2.2]). Hence the automorphism α stated in the lemma above is also asymptotically representable. Besides, it is not so hard to see that the crossed product C^* -algebra $A \rtimes_\sigma \mathbb{Z}$ is a unital simple AT algebra with real rank zero. Therefore, $A \rtimes_\alpha \mathbb{Z}$ is a unital simple AT algebra with real rank zero, too.

5 The Rohlin property of $\mathbb{Z}^2$ -actions on AF algebras

In this section, we would like to show that certain $\mathbb{Z}^2$ -actions on an AF algebra have the Rohlin property. This is a generalization of Nakamura's theorem [23, Theorem 3].

Throughout this section, we keep the following setting. Let A be a unital simple separable C^* -algebra with tracial rank zero and suppose that A has a unique tracial state τ . Suppose that automorphisms $\alpha, \beta \in \text{Aut}(A)$ and a unitary $w \in A$ satisfy

$$\beta \circ \alpha = \text{Ad } w \circ \alpha \circ \beta$$

and $\alpha^m \circ \beta^n$ is uniformly outer for all $(m, n) \in \mathbb{Z}^2 \setminus \{(0, 0)\}$. We remark that α and β induce a $\mathbb{Z}^2$ -action on A_∞ .

Lemma 5.1. *For any $m_1, m_2 \in \mathbb{N}$, there exists a central sequence of projections $(e_n)_n$ in A such that*

$$\lim_{n \rightarrow \infty} \tau(e_n) = \frac{1}{m_1 m_2}$$

and

$$\lim_{n \rightarrow \infty} \|\beta^j(\alpha^i(e_n))\beta^l(\alpha^k(e_n))\| = 0$$

for all $(i, j) \neq (k, l)$ in $\{(i, j) \mid 0 \leq i \leq m_1 - 1, 0 \leq j \leq m_2 - 1\}$.

Proof. Set $I = \{(i, j) \mid 0 \leq i \leq m_1 - 1, 0 \leq j \leq m_2 - 1\}$. Let (π_τ, H_τ) be the GNS representation associated with τ . It is well-known that $\pi_\tau(A)''$ is a hyperfinite II_1 -factor (see [26, Lemma 2.16]). For $(i, j) \in \mathbb{Z}^2$, we put $\varphi_{(i, j)} = \alpha^i \circ \beta^j$. Then $\varphi : \mathbb{Z}^2 \rightarrow \text{Aut}(A)$ is a cocycle action of $\mathbb{Z}^2$. We denote its extension to $\pi_\tau(A)''$ by $\bar{\varphi}$. Since $\alpha^m \circ \beta^n$ is uniformly outer for all $(m, n) \in \mathbb{Z}^2 \setminus \{(0, 0)\}$, $\bar{\varphi}$ is an outer cocycle action of $\mathbb{Z}^2$ on $\pi_\tau(A)''$. It follows from [25] that there exists a sequence of projections $(e_n)_n$ in $\pi_\tau(A)''$ such that

$$\sum_{(i, j) \in I} \beta^j(\alpha^i(e_n)) \rightarrow 1, \quad [x, e_n] \rightarrow 0 \quad \forall x \in \pi_\tau(A)''$$

and

$$\beta^j(\alpha^i(e_n))\beta^l(\alpha^k(e_n)) \rightarrow 0 \quad \forall (i, j), (k, l) \in I \text{ with } (i, j) \neq (k, l)$$

in the strong operator topology as $n \rightarrow \infty$. By [26, Lemma 2.15], we may replace e_n with projections in A . Applying Proposition 4.1 to $\Gamma = \{\beta^j \circ \alpha^i \mid (i, j) \in I\}$, we obtain the conclusion. $\square$

Let $(e_n)_n$ be the projections as in the lemma above. If α^r is in $\overline{\text{Inn}}(A)$ for some $r \in \mathbb{N}$ and m_1 is large enough, then we can construct a central sequence of projections $(e'_n)_n$ in A such that

$$\lim_{n \rightarrow \infty} \|e'_n(e_n + \alpha(e_n) + \cdots + \alpha^{m_1-1}(e_n)) - e'_n\| = 0,$$

$$\alpha(e'_n) \approx e'_n \quad \text{and} \quad \tau(e'_n) \approx m_1 \tau(e_n),$$

by using the arguments in [9, Lemma 3.1] (see also [23, Lemma 6]). Consequently, we get the following.

Lemma 5.2. *If α^r is in $\overline{\text{Inn}}(A)$ for some $r \in \mathbb{N}$, then for any $m \in \mathbb{N}$, there exists a central sequence of projections $(e_n)_n$ in A such that*

$$\lim_{n \rightarrow \infty} \tau(e_n) = \frac{1}{m}, \quad \lim_{n \rightarrow \infty} \|e_n - \alpha(e_n)\| = 0$$

and

$$\lim_{n \rightarrow \infty} \|e_n \beta^j(e_n)\| = 0$$

for all $j = 1, 2, \dots, m-1$.

Our next task is to achieve the cyclicity condition $\beta^m(e_n) \approx e_n$.

Lemma 5.3. *Suppose that A is AF. Suppose that a projection $e \in A_\infty$ and a partial isometry $u \in A_\infty$ satisfy $e = \alpha(e)$ and $e = u^*u = uu^*$. Then there exists a partial isometry $w \in A_\infty$ such that $w^*w = ww^* = e$ and $u = w^*\alpha(w)$.*

Proof. Theorem 4.8 tells us that α possesses the Rohlin property. We can modify the standard argument deducing stability from the Rohlin property (see [6, 4]) and apply it to the unitary $u + (1 - e)$. We leave the details to the readers. $\square$

Lemma 5.4. *Suppose that either of the following holds.*

- (1) *A is AF, α^r is approximately inner for some $r \in \mathbb{N}$ and β^s is approximately inner for some $s \in \mathbb{N}$.*
- (2) *α^r is approximately inner for some $r \in \mathbb{N}$ and there exist a natural number $s \in \mathbb{N}$ and a sequence of unitaries $(u_n)_n$ in A such that*

$$\lim_{n \rightarrow \infty} \|u_n - \alpha(u_n)\| = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \|u_n \alpha u_n^* - \beta^s(a)\| = 0 \quad \forall a \in A.$$

Then for any $m \in \mathbb{N}$, there exists a central sequence of projections $(e_n)_n$ such that

$$\lim_{n \rightarrow \infty} \tau(e_n) = \frac{1}{m}, \quad \lim_{n \rightarrow \infty} \|e_n - \alpha(e_n)\| = 0, \quad \lim_{n \rightarrow \infty} \|e_n \beta^j(e_n)\| = 0$$

for all $j = 1, 2, \dots, m-1$ and

$$\lim_{n \rightarrow \infty} \|e_n - \beta^m(e_n)\| = 0.$$

Proof. Choose a large natural number l such that $l \equiv 1 \pmod{s}$. By using Lemma 5.2 and the assumption that β^s is in $\overline{\text{Inn}}(A)$ for some $s \in \mathbb{N}$, one can find a projection $e \in A_\infty$ and a partial isometry $v \in A_\infty$ such that

$$e = \alpha(e), \quad v^*v = e, \quad vv^* = \beta(e) \quad \text{and} \quad e\beta^j(e) = 0 \quad \forall j = 1, 2, \dots, l-1$$

in the same way as the proof of Theorem 4.4. Moreover, we have

$$\lim_{n \rightarrow \infty} \tau(e_n) = l^{-1},$$

where $(e_n)_n$ is a representative sequence of e consisting of projections. Note that $\beta^j(e)$ is fixed by α , because e is a central sequence. In the case (2), clearly we may further assume $v = \alpha(v)$. In the case (1), the lemma above applies to $v^*\alpha(v)$ and yields $w \in A_\infty$ satisfying $w^*w = e$, $ww^* = e$ and $v^*\alpha(v) = w^*\alpha(w)$. By replacing v with vw^* , we get $v = \alpha(v)$, too. Then the conclusion follows from exactly the same argument as [9, Lemma 4.3]. $\square$

Theorem 5.5. *Suppose that the conclusion of Lemma 5.4 holds. Then for any $m \in \mathbb{N}$, there exist projections e and f in A_∞ such that*

$$\alpha(e) = e, \quad \alpha(f) = f, \quad \beta^m(e) = e, \quad \beta^{m+1}(f) = f$$

and

$$\sum_{i=0}^{m-1} \beta^i(e) + \sum_{j=0}^m \beta^j(f) = 1.$$

Proof. Let $(e_n)_n$ be the central sequence of projections obtained in Lemma 5.4. Define

$$f_n = 1 - \sum_{j=0}^{m-1} \beta^j(e_n).$$

There exists a sequence of unitaries $(u_n)_n$ in A such that $u_n \rightarrow 1$ as $n \rightarrow \infty$ and $u_n \alpha(e_n) u_n^* = e_n$ for sufficiently large n . The $\mathbb{Z}$ -action on $e_n A e_n$ generated by $\text{Ad } u_n \circ \alpha$ is uniformly outer, and so it has the tracial Rohlin property by Theorem 4.2 (or [26, Theorem 2.17]). It follows that, for any $k \in \mathbb{N}$, there exists a central sequence of projections $(\tilde{e}_n)_n$ such that

$$\tilde{e}_n \leq e_n, \quad \lim_{n \rightarrow \infty} \tau(\tilde{e}_n) = 1/mk, \quad \text{and} \quad \lim_{n \rightarrow \infty} \|\tilde{e}_n \alpha^i(\tilde{e}_n)\| = 0 \quad \forall i = 1, 2, \dots, k-1.$$

Let $e, f, \tilde{e} \in A_\infty$ be the images of $(e_n)_n, (f_n)_n, (\tilde{e}_n)_n$, respectively. By Lemma 3.3, there exists a partial isometry v such that $v^*v = f$ and $vv^* \leq \tilde{e}$. We define a partial isometry $\tilde{v} \in A_\infty$ by

$$\tilde{v} = \frac{1}{\sqrt{k}} \sum_{i=0}^{k-1} \alpha^i(v).$$

Then one has

$$\tilde{v}^* \tilde{v} = f, \quad \tilde{v} \tilde{v}^* \leq e \quad \text{and} \quad \|\tilde{v} - \alpha(\tilde{v})\| < 2/\sqrt{k}.$$

By a standard trick on central sequences, we may assume $\alpha(\tilde{v}) = \tilde{v}$. Thus, we have obtained the α -invariant version of the conclusion of Lemma 4.6. We can complete the proof by the same argument as in Theorem 4.7. $\square$

The following is a generalization of [23, Theorem 3].

Corollary 5.6. *Let $\varphi : \mathbb{Z}^2 \curvearrowright A$ be a $\mathbb{Z}^2$ -action on a unital simple AF algebra A with unique trace. When $\varphi_{(r,0)}$ and $\varphi_{(0,s)}$ are approximately inner for some $r, s \in \mathbb{N}$, the following are equivalent.*

- (1) φ has the Rohlin property.
- (2) φ is uniformly outer.

Proof. This immediately follows from Theorem 5.5 and [23, Remark 2] (see also [22, Remark 2.2]). $\square$

The next corollary also follows from Theorem 5.5 immediately, because condition (2) of Lemma 5.4 is satisfied in this case. See [7, Definition 2.2] for the definition of approximate representability.

Corollary 5.7. *Let $\varphi : \mathbb{Z}^2 \curvearrowright A$ be an approximately representable $\mathbb{Z}^2$ -action on a unital simple AH algebra A with real rank zero and slow dimension growth. Suppose that A has a unique trace. Then the following are equivalent.*

- (1) φ has the Rohlin property.
- (2) φ is uniformly outer.

6 Classification of certain $\mathbb{Z}^2$ -actions on AF algebras

In this section, we will show a classification result of a certain class of $\mathbb{Z}^2$ -actions on unital simple AF algebras. We freely use the terminology and notation introduced in [7, Definition 2.1]. For an automorphism α of a C^* -algebra A , we write the crossed product C^* -algebra $A \rtimes_{\alpha} \mathbb{Z}$ by $C^*(A, \alpha)$ and the implementing unitary by λ_{α} . The mapping torus $M(A, \alpha)$ is defined by

$$M(A, \alpha) = \{f \in C([0, 1], A) \mid \alpha(f(0)) = f(1)\}.$$

When A is an AF algebra, ‘ KK -triviality’ of $\alpha \in \text{Aut}(A)$ is equivalent to $K_0(\alpha) = \text{id}$, and also equivalent to α being approximately inner.

The following theorem is a $\mathbb{Z}$ -equivariant version of Theorem 4.9. Let A be a unital simple AF algebra with unique trace and let $\alpha \in \overline{\text{Inn}}(A)$. Let $\text{Aut}_{\mathbb{T}}(C^*(A, \alpha))$ denote the set of all automorphisms of $C^*(A, \alpha)$ commuting with the dual action $\hat{\alpha}$. For $i = 1, 2$, we suppose that an automorphism $\beta_i \in \text{Aut}(A)$ and a unitary $w_i \in A$ are given and satisfy

$$\beta_i \circ \alpha = \text{Ad } w_i \circ \alpha \circ \beta_i.$$

Then β_i extends to $\tilde{\beta}_i \in \text{Aut}_{\mathbb{T}}(C^*(A, \alpha))$ by setting $\tilde{\beta}_i(\lambda_{\alpha}) = w_i \lambda_{\alpha}$. Suppose further that $\alpha^m \circ \beta_i^n$ is uniformly outer for all $(m, n) \in \mathbb{Z}^2 \setminus \{(0, 0)\}$ and that $\beta_i^{s_i}$ is approximately inner for some $s_i \in \mathbb{N}$.

Theorem 6.1. *In the setting above, if $\tilde{\beta}_1$ and $\tilde{\beta}_2$ are asymptotically unitarily equivalent, then there exist an approximately inner automorphism $\mu \in \text{Aut}_{\mathbb{T}}(C^*(A, \alpha))$ and a unitary $v \in A$ such that $\mu|_A$ is also approximately inner and*

$$\mu \circ \tilde{\beta}_1 \circ \mu^{-1} = \text{Ad } v \circ \tilde{\beta}_2.$$

Proof. We can apply the argument of [24, Theorem 5] to $\tilde{\beta}_1$ and $\tilde{\beta}_2$ in a similar fashion to [7, Theorem 4.11]. By Remark 4.11, (the $\mathbb{Z}$ -action generated by) α is asymptotically representable. Then [7, Theorem 4.8] implies that $\tilde{\beta}_1$ and $\tilde{\beta}_2$ are $\mathbb{T}$ -asymptotically unitarily equivalent. Moreover, by Theorem 5.5, we can find Rohlin projections for $\tilde{\beta}_i$ in the fixed point algebra $(A_{\infty})^{\alpha}$. Hence, by using Lemma 4.10 instead of [24, Theorem 7], the usual intertwining argument shows the statement. $\square$

Let us recall the OrderExt invariant introduced in [13]. Let G_0, G_1, F be abelian groups and let $D : G_0 \rightarrow F$ be a homomorphism. When

$$\xi : \quad 0 \longrightarrow G_0 \xrightarrow{\iota} E_{\xi} \xrightarrow{q} G_1 \longrightarrow 0$$

is exact, R is in $\text{Hom}(E_\xi, F)$ and $R \circ \iota = D$, the pair (ξ, R) is called an order-extension. Two order-extensions (ξ, R) and (ξ', R') are equivalent if there exists an isomorphism $\theta : E_\xi \rightarrow E_{\xi'}$ such that $R = R' \circ \theta$ and

$$\begin{array}{ccccccccc} \xi : & 0 & \longrightarrow & G_0 & \longrightarrow & E_\xi & \longrightarrow & G_1 & \longrightarrow & 0 \\ & & & \parallel & & \downarrow \theta & & \parallel & & \\ \xi' : & 0 & \longrightarrow & G_0 & \longrightarrow & E_{\xi'} & \longrightarrow & G_1 & \longrightarrow & 0 \end{array}$$

is commutative. Then $\text{OrderExt}(G_1, G_0, D)$ consists of equivalence classes of all order-extensions. As shown in [13], $\text{OrderExt}(G_1, G_0, D)$ is equipped with an abelian group structure. The map sending (ξ, R) to ξ induces a homomorphism from $\text{OrderExt}(G_1, G_0, D)$ onto $\text{Ext}(G_1, G_0)$.

Let B be a unital C^* -algebra with $T(B)$ non-empty. We denote by $\text{Aut}_0(B)$ the set of all automorphisms γ of B such that $K_0(\gamma) = K_1(\gamma) = \text{id}$ and $\tau \circ \gamma = \tau$ for all $\tau \in T(B)$. When B is a unital simple AT algebra with real rank zero, $\text{Aut}_0(B)$ equals $\overline{\text{Inn}}(B)$. Let $D_B : K_0(B) \rightarrow \text{Aff}(T(B))$ denote the dimension map defined by $D_B([p])(\tau) = \tau(p)$. As described in [13], there exist natural homomorphisms

$$\tilde{\eta}_0 : \text{Aut}_0(B) \rightarrow \text{OrderExt}(K_1(B), K_0(B), D_B)$$

and

$$\eta_1 : \text{Aut}_0(B) \rightarrow \text{Ext}(K_0(B), K_1(B)).$$

The following is the main result of [13]. See [21, 18] for further developments.

Theorem 6.2 ([13, Theorem 4.4]). *Suppose that B is a unital simple AT algebra with real rank zero. Then the homomorphism*

$$\tilde{\eta}_0 \oplus \eta_1 : \overline{\text{Inn}}(B) \rightarrow \text{OrderExt}(K_1(B), K_0(B), D_B) \oplus \text{Ext}(K_0(B), K_1(B))$$

is surjective and its kernel equals the set of all asymptotically inner automorphisms of B .

By using this OrderExt invariant, we introduce an invariant of certain $\mathbb{Z}^2$ -actions as follows. Let A be a unital simple AF algebra and let $\varphi : \mathbb{Z}^2 \curvearrowright A$ be an action of $\mathbb{Z}^2$ on A . Suppose that φ is uniformly outer and locally KK -trivial (i.e. locally approximately inner). We write $B = C^*(A, \varphi_{(1,0)})$. Then $\varphi_{(0,1)}$ extends to $\tilde{\varphi}_{(0,1)} \in \text{Aut}(B)$ by setting $\tilde{\varphi}_{(0,1)}(\lambda_{\varphi_{(1,0)}}) = \lambda_{\varphi_{(1,0)}}$. Let $\iota : A \rightarrow B = C^*(A, \varphi_{(1,0)})$ be the canonical inclusion. One can check the following immediately.

- $K_0(\iota)$ is an isomorphism from $K_0(A)$ to $K_0(B)$.
- The connecting map $\partial : K_1(B) \rightarrow K_0(A)$ in the Pimsner-Voiculescu exact sequence is an isomorphism and $\partial^{-1}([p]) = [\lambda_{\varphi_{(1,0)}} \iota(p) + \iota(v(1-p))]$ for any projection $p \in A$, where v is a unitary of A satisfying $vpv^* = \varphi_{(1,0)}(p)$.
- The map $\iota^* : T(B) \rightarrow T(A)$ sending τ to $\tau \circ \iota$ is an isomorphism and satisfies $D_B(K_0(\iota)(x))(\tau) = D_A(x)(\iota^*(\tau))$ for $x \in K_0(A)$ and $\tau \in T(B)$.

From these properties, we can obtain a natural isomorphism

$$\zeta_{\varphi_{(1,0)}} : \text{OrderExt}(K_1(B), K_0(B), D_B) \rightarrow \text{OrderExt}(K_0(A), K_0(A), D_A).$$

In addition, it is easy to see $K_0(\tilde{\varphi}_{(0,1)}) = K_1(\tilde{\varphi}_{(0,1)}) = \text{id}$ and $\tau \circ \tilde{\varphi}_{(0,1)} = \tau$ for all $\tau \in T(B)$, that is, $\tilde{\varphi}_{(0,1)}$ belongs to $\text{Aut}_0(B)$.

Lemma 6.3. *In the setting above, $\eta_1(\tilde{\varphi}_{(0,1)}) \in \text{Ext}(K_0(B), K_1(B))$ is zero.*

Proof. There exists a natural commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & C_0((0,1), B) & \longrightarrow & M(B, \tilde{\varphi}_{(0,1)}) & \longrightarrow & B & \longrightarrow & 0 \\ & & \uparrow & & \uparrow & & \uparrow \iota & & \\ 0 & \longrightarrow & C_0((0,1), A) & \longrightarrow & M(A, \varphi_{(0,1)}) & \longrightarrow & A & \longrightarrow & 0, \end{array}$$

where the horizontal sequences are exact. From the naturality of the six-term exact sequence, we obtain the commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & K_1(B) & \longrightarrow & K_0(M(B, \tilde{\varphi}_{(0,1)})) & \longrightarrow & K_0(B) & \longrightarrow & 0 \\ & & \uparrow & & \uparrow & & \uparrow K_0(\iota) & & \\ 0 & \longrightarrow & K_1(A) & \longrightarrow & K_0(M(A, \varphi_{(0,1)})) & \longrightarrow & K_0(A) & \longrightarrow & 0, \end{array}$$

where the horizontal sequences are exact. Since $K_1(A)$ is zero and $K_0(\iota)$ is an isomorphism, we can conclude $\eta_1(\tilde{\varphi}_{(0,1)}) = 0$. $\square$

Definition 6.4. In the setting above, we define our invariant $[\varphi]$ by

$$[\varphi] = \zeta_{\varphi_{(1,0)}}(\tilde{\eta}_0(\tilde{\varphi}_{(0,1)})) \in \text{OrderExt}(K_0(A), K_0(A), D_A).$$

Proposition 6.5. *Let $\varphi, \psi : \mathbb{Z}^2 \curvearrowright A$ be uniformly outer, locally KK -trivial $\mathbb{Z}^2$ -actions on a unital simple AF algebra A . If φ and ψ are KK -trivially cocycle conjugate, then $[\varphi] = [\psi]$.*

Proof. For $\mu \in \overline{\text{Inn}}(A)$, it is straightforward to see that the $\mathbb{Z}^2$ -action $\mu \circ \varphi \circ \mu^{-1}$ has the same invariant as φ . Hence, it suffices to show $[\varphi] = [\varphi^u]$ for any φ -cocycle $\{u_n\}_{n \in \mathbb{Z}^2}$. Define an isomorphism π from $C^*(A, \varphi_{(1,0)})$ to $C^*(A, \varphi_{(1,0)}^u)$ by

$$\pi(\lambda_{\varphi_{(1,0)}}) = u_{(1,0)}^* \lambda_{\varphi_{(1,0)}^u} \quad \text{and} \quad \pi(a) = a \quad \forall a \in A,$$

where A is identified with subalgebras of the crossed products. For $\gamma \in \text{Aut}(C^*(A, \varphi_{(1,0)}^u))$, one can check

$$\zeta_{\varphi_{(1,0)}}(\tilde{\eta}_0(\pi^{-1} \circ \gamma \circ \pi)) = \zeta_{\varphi_{(1,0)}^u}(\tilde{\eta}_0(\gamma)) \in \text{OrderExt}(K_0(A), K_0(A), D_A),$$

where $\tilde{\eta}_0$ in the left hand side is defined for $C^*(A, \varphi_{(1,0)})$ and $\tilde{\eta}_0$ in the right hand side is defined for $C^*(A, \varphi_{(1,0)}^u)$. We also have

$$(\pi^{-1} \circ \tilde{\varphi}_{(0,1)}^u \circ \pi)(a) = \pi^{-1}(\tilde{\varphi}_{(0,1)}^u(a)) = \tilde{\varphi}_{(0,1)}^u(a) = (\text{Ad } u_{(0,1)} \circ \tilde{\varphi}_{(0,1)})(a) \quad \forall a \in A$$

and

$$\begin{aligned}
(\pi^{-1} \circ \tilde{\varphi}_{(0,1)}^u \circ \pi)(\lambda_{\varphi_{(1,0)}}) &= (\pi^{-1} \circ \tilde{\varphi}_{(0,1)}^u)(u_{(1,0)}^* \lambda_{\varphi_{(1,0)}^u}) \\
&= \pi^{-1}(\tilde{\varphi}_{(0,1)}^u(u_{(1,0)}^*) \lambda_{\varphi_{(1,0)}^u}) \\
&= \varphi_{(0,1)}^u(u_{(1,0)}^*) u_{(1,0)} \lambda_{\varphi_{(1,0)}} \\
&= u_{(0,1)} \varphi_{(0,1)}(u_{(1,0)}^*) u_{(0,1)}^* u_{(1,0)} \lambda_{\varphi_{(1,0)}} \\
&= u_{(0,1)} \varphi_{(1,0)}(u_{(0,1)}^*) u_{(1,0)}^* u_{(1,0)} \lambda_{\varphi_{(1,0)}} \\
&= u_{(0,1)} \varphi_{(1,0)}(u_{(0,1)}^*) \lambda_{\varphi_{(1,0)}} \\
&= u_{(0,1)} \lambda_{\varphi_{(1,0)}} u_{(0,1)}^* \\
&= (\text{Ad } u_{(0,1)} \circ \tilde{\varphi}_{(0,1)})(\lambda_{\varphi_{(1,0)}}).
\end{aligned}$$

Thus $\pi^{-1} \circ \tilde{\varphi}_{(0,1)}^u \circ \pi = \text{Ad } u_{(0,1)} \circ \tilde{\varphi}_{(0,1)}$. Since inner automorphisms are contained in the kernel of $\tilde{\eta}_0$, we obtain

$$\begin{aligned}
\zeta_{\varphi_{(1,0)}}(\tilde{\eta}_0(\tilde{\varphi}_{(0,1)})) &= \zeta_{\varphi_{(1,0)}}(\tilde{\eta}_0(\text{Ad } u_{(0,1)} \circ \tilde{\varphi}_{(0,1)})) \\
&= \zeta_{\varphi_{(1,0)}}(\tilde{\eta}_0(\pi^{-1} \circ \tilde{\varphi}_{(0,1)}^u \circ \pi)) \\
&= \zeta_{\varphi_{(1,0)}^u}(\tilde{\eta}_0(\tilde{\varphi}_{(0,1)}^u)),
\end{aligned}$$

which completes the proof. $\square$

Theorem 6.6. *Let $\varphi, \psi : \mathbb{Z}^2 \curvearrowright A$ be uniformly outer, locally KK -trivial $\mathbb{Z}^2$ -actions on a unital simple AF algebra A with unique trace. The following are equivalent.*

- (1) $[\varphi] = [\psi]$.
- (2) φ and ψ are KK -trivially cocycle conjugate.

Proof. (2) $\Rightarrow$ (1) was shown in the proposition above without assuming that A has a unique trace. Let us consider the other implication (1) $\Rightarrow$ (2). By Theorem 4.8 and Theorem 4.9, we may assume that there exists a unitary $u \in A$ such that $\psi_{(1,0)} = \text{Ad } u \circ \varphi_{(1,0)}$. By Theorem 4.8 and Remark 4.11 (or Remark 4.5), the crossed product C^* -algebra $C^*(A, \varphi_{(1,0)})$ is a unital simple AT algebra with real rank zero.

Clearly $\varphi_{(0,1)}$ extends to $\tilde{\varphi}_{(0,1)} \in \text{Aut}(C^*(A, \varphi_{(1,0)}))$ by

$$\tilde{\varphi}_{(0,1)}(a) = a \quad \forall a \in A \quad \text{and} \quad \tilde{\varphi}_{(0,1)}(\lambda_{\varphi_{(1,0)}}) = \lambda_{\varphi_{(1,0)}}.$$

Since

$$\psi_{(0,1)} \circ \varphi_{(1,0)} = \text{Ad}(\psi_{(0,1)}(u^*)u) \circ \varphi_{(1,0)} \circ \psi_{(0,1)},$$

we can extend $\psi_{(0,1)}$ to $\omega \in \text{Aut}(C^*(A, \varphi_{(1,0)}))$ by

$$\omega(a) = a \quad \forall a \in A \quad \text{and} \quad \omega(\lambda_{\varphi_{(1,0)}}) = \psi_{(0,1)}(u^*)u \lambda_{\varphi_{(1,0)}}.$$

In order to apply Theorem 6.1 to $\tilde{\varphi}_{(0,1)}$ and ω , we would like to check that these automorphisms are asymptotically unitarily equivalent. There exists an isomorphism $\pi : C^*(A, \varphi_{(1,0)}) \rightarrow C^*(A, \psi_{(1,0)})$ defined by

$$\pi(a) = a \quad \forall a \in A \quad \text{and} \quad \pi(\lambda_{\varphi_{(1,0)}}) = u^* \lambda_{\psi_{(1,0)}}.$$

As mentioned in the proof of Proposition 6.5, for any $\gamma \in \text{Aut}(C^*(A, \varphi_{(1,0)}))$, one has

$$\zeta_{\varphi_{(1,0)}}(\tilde{\eta}_0(\gamma)) = \zeta_{\psi_{(1,0)}}(\tilde{\eta}_0(\pi \circ \gamma \circ \pi^{-1})).$$

Moreover it is easy to see that $\pi \circ \omega \circ \pi^{-1}$ is equal to $\tilde{\psi}_{(0,1)}$, which is defined by

$$\tilde{\psi}_{(0,1)}(a) = a \quad \forall a \in A \quad \text{and} \quad \tilde{\psi}_{(0,1)}(\lambda_{\psi_{(1,0)}}) = \lambda_{\psi_{(1,0)}}.$$

It follows that

$$\begin{aligned} \zeta_{\varphi_{(1,0)}}(\tilde{\eta}_0(\omega)) &= \zeta_{\psi_{(1,0)}}(\tilde{\eta}_0(\pi \circ \omega \circ \pi^{-1})) \\ &= \zeta_{\psi_{(1,0)}}(\tilde{\eta}_0(\tilde{\psi}_{(0,1)})) = [\psi] = [\varphi] = \zeta_{\varphi_{(1,0)}}(\tilde{\eta}_0(\tilde{\varphi}_{(0,1)})), \end{aligned}$$

and so $\tilde{\eta}_0(\omega) = \tilde{\eta}_0(\tilde{\varphi}_{(0,1)})$. By Lemma 6.3, $\eta_1(\omega) = \eta_1(\tilde{\varphi}_{(0,1)}) = 0$. Therefore, by Theorem 6.2, $\tilde{\varphi}_{(0,1)}$ and ω are asymptotically unitarily equivalent.

Then, Theorem 6.1 applies and yields an approximately inner automorphism $\mu \in \text{Aut}_{\mathbb{T}}(C^*(A, \varphi_{(1,0)}))$ and a unitary $v \in A$ such that $\mu|_A$ is in $\overline{\text{Inn}}(A)$ and

$$\mu \circ \omega \circ \mu^{-1} = \text{Ad } v \circ \tilde{\varphi}_{(0,1)}. \quad (6.1)$$

By restricting this equality to A , we get

$$(\mu|_A) \circ \psi_{(0,1)} \circ (\mu|_A)^{-1} = \text{Ad } v \circ \varphi_{(0,1)}. \quad (6.2)$$

Let $z \in A$ be the unitary satisfying $\mu(\lambda_{\varphi_{(1,0)}}) = z\lambda_{\varphi_{(1,0)}}$. Then

$$(\mu|_A) \circ \psi_{(1,0)} \circ (\mu|_A)^{-1} = (\mu|_A) \circ \text{Ad } u \circ \varphi_{(1,0)} \circ (\mu|_A)^{-1} = \text{Ad } \mu(u)z \circ \varphi_{(1,0)}. \quad (6.3)$$

From (6.1), one can see that

$$\begin{aligned} (\mu \circ \omega \circ \mu^{-1})(\lambda_{\varphi_{(1,0)}}) &= (\mu \circ \omega)(\mu^{-1}(z^*)\lambda_{\varphi_{(1,0)}}) \\ &= \mu(\psi_{(0,1)}(\mu^{-1}(z^*))\psi_{(0,1)}(u^*)u\lambda_{\varphi_{(1,0)}}) \\ &= (\text{Ad } v \circ \varphi_{(0,1)})(z^*\mu(u^*))\mu(u)z\lambda_{\varphi_{(1,0)}} \\ &= v\varphi_{(0,1)}(z^*\mu(u^*))v^*\mu(u)z\lambda_{\varphi_{(1,0)}} \end{aligned}$$

is equal to

$$\begin{aligned} (\text{Ad } v \circ \tilde{\varphi}_{(0,1)})(\lambda_{\varphi_{(1,0)}}) &= v\lambda_{\varphi_{(1,0)}}v^* \\ &= v\varphi_{(1,0)}(v^*)\lambda_{\varphi_{(1,0)}}. \end{aligned}$$

Hence one obtains

$$v\varphi_{(0,1)}(\mu(u)z) = \mu(u)z\varphi_{(1,0)}(v). \quad (6.4)$$

It follows from (6.2), (6.3), (6.4) that ψ and φ are KK -trivially cocycle conjugate. $\square$

Remark 6.7. We do not know the precise range of our invariant which takes its values in OrderExt . At least, the following observation shows that the range does not exhaust OrderExt . Let $\varphi : \mathbb{Z}^2 \curvearrowright A$ be a locally KK -trivial and uniformly outer $\mathbb{Z}^2$ -action on a unital simple AF algebra. Suppose that (ξ, R) is a representative of $[\varphi] \in \text{OrderExt}(K_0(A), K_0(A), D_A)$. Since

$$\xi : 0 \longrightarrow K_0(A) \xrightarrow{\iota} E_\xi \xrightarrow{q} K_0(A) \longrightarrow 0$$

is exact and $R : E_\xi \rightarrow \text{Aff}(T(A))$ satisfies $R \circ \iota = D_A$, there exists a homomorphism $R_0 : K_0(A) \rightarrow \text{Aff}(K_0(A))/\text{Im } D_A$ such that $R_0(q(x)) = R(x) + D_A(K_0(A))$ for any $x \in E_\xi$. It is easy to see $R_0([1_A]) = 0$, because the implementing unitary $\lambda_{\varphi(1,0)}$ is fixed by $\tilde{\varphi}_{(0,1)}$. Thus, $[\varphi]$ belongs to the subgroup

$$\{[(\xi, R)] \in \text{OrderExt}(K_0(A), K_0(A), D_A) \mid R_0([1_A]) = 0\}.$$

When A is a UHF algebra, one can see that this subgroup coincides with the range of the invariant introduced in [8]. Therefore, Theorem 6.6 yields a new proof of [8, Theorem 6.5].

References

- [1] M. Dadarlat, *Reduction to dimension three of local spectra of real rank zero C^* -algebras*, J. Reine Angew. Math. 460 (1995), 189–212.
- [2] G. A. Elliott and G. Gong, *On the classification of C^* -algebras of real rank zero. II*, Ann. of Math. (2) 144 (1996), 497–610.
- [3] G. A. Elliott, G. Gong and L. Li, *Approximate divisibility of simple inductive limit C^* -algebras*, Operator algebras and operator theory (Shanghai, 1997), 87–97, Contemp. Math. 228, Amer. Math. Soc., Providence, RI, 1998.
- [4] D. E. Evans and A. Kishimoto, *Trace scaling automorphisms of certain stable AF algebras*, Hokkaido Math. J. 26 (1997), 211–224.
- [5] G. Gong, *On inductive limits of matrix algebras over higher-dimensional spaces. II*, Math. Scand. 80 (1997), 56–100.
- [6] R. H. Herman and A. Ocneanu, *Stability for integer actions on UHF C^* -algebras*, J. Funct. Anal. 59 (1984), 132–144.
- [7] M. Izumi and H. Matui, *$\mathbb{Z}^2$ -actions on Kirchberg algebras*, preprint. arXiv:0902.0194
- [8] T. Katsura and H. Matui, *Classification of uniformly outer actions of $\mathbb{Z}^2$ on UHF algebras*, Adv. Math. 218 (2008), 940–968.
- [9] A. Kishimoto, *The Rohlin property for automorphisms of UHF algebras*, J. Reine Angew. Math. 465 (1995), 183–196.
- [10] A. Kishimoto, *The Rohlin property for shifts on UHF algebras and automorphisms of Cuntz algebras*, J. Funct. Anal. 140 (1996), 100–123.

- [11] A. Kishimoto, *Automorphisms of AT algebras with the Rohlin property*, J. Operator Theory 40 (1998), 277–294.
- [12] A. Kishimoto, *Unbounded derivations in AT algebras*, J. Funct. Anal. 160 (1998), 270–311.
- [13] A. Kishimoto and A. Kumjian, *The Ext class of an approximately inner automorphism, II*, J. Operator Theory, 46 (2001), 99–122.
- [14] H. Lin, *Tracially AF C^* -algebras*, Trans. Amer. Math. Soc. 353 (2001), 693–722.
- [15] H. Lin, *The tracial topological rank of C^* -algebras*, Proc. London Math. Soc. 83 (2001), 199–234.
- [16] H. Lin, *Classification of simple C^* -algebras with tracial topological rank zero*, Duke Math. J. 125 (2004), 91–119.
- [17] H. Lin, *The Rokhlin property for automorphisms on simple C^* -algebras*, Operator theory, operator algebras, and applications, 189–215, Contemp. Math. 414, Amer. Math. Soc., Providence, RI, 2006.
- [18] H. Lin, *Asymptotically unitary equivalence and asymptotically inner automorphisms*, preprint. arXiv:math/0703610
- [19] H. Lin and H. Matui, *Minimal dynamical systems on the product of the Cantor set and the circle II*, Selecta Math. (N.S.) 12 (2006), 199–239.
- [20] H. Lin and H. Osaka, *The Rokhlin property and the tracial topological rank*, J. Funct. Anal. 218 (2005), 475–494.
- [21] H. Matui, *AF embeddability of crossed products of AT algebras by the integers and its application*, J. Funct. Anal. 192 (2002), 562–580.
- [22] H. Matui, *Classification of outer actions of $\mathbb{Z}^N$ on $\mathcal{O}_2$* , Adv. Math. 217 (2008), 2872–2896.
- [23] H. Nakamura, *The Rohlin property for $\mathbb{Z}^2$ -actions on UHF algebras*, J. Math. Soc. Japan 51 (1999), 583–612.
- [24] H. Nakamura, *Aperiodic automorphisms of nuclear purely infinite simple C^* -algebras*, Ergodic Theory Dynam. Systems 20 (2000), 1749–1765.
- [25] A. Ocneanu, *Actions of discrete amenable groups on von Neumann algebras*, Lecture Notes in Mathematics 1138, Springer-Verlag, Berlin, 1985.
- [26] H. Osaka and N. C. Phillips, *Furstenberg transformations on irrational rotation algebras*, Ergodic Theory Dynam. Systems 26 (2006), 1623–1651.
- [27] Y. Sato, *Certain aperiodic automorphisms of unital simple projectionless C^* -algebras*, to appear in Internat. J. Math.